
PART 5


LM FOOD CHAIN

Appendix 5.5.2. Agreement Between Modeled and Observed PCB Concentrations


Agreement Between Modeled and Observed PCB Concentrations in Zooplankton, *Mysis*, and *Diporeia* at Sturgeon Bay (1994 and 1995).




Agreement Between Modeled and Observed Fish PCB Concentrations in Forage Fish at Sturgeon Bay (1994 and 1995)




Agreement Between Modeled and Observed Fish PCB Concentrations in Lake Trout at Sturgeon Bay (1994 and 1995).




Agreement Between Modeled and Observed PCB Concentrations in Zooplankton, *Mysis*, and *Diporeia* at Sheboygan Reef (1994 and 1995)




Agreement Between Modeled and Observed Fish PCB Concentrations in Forage Fish at Sheboygan Reef (1994 and 1995)




Agreement Between Modeled and Observed Fish PCB Concentrations in Lake Trout at Sheboygan Reef (1994 and 1995)



Agreement Between Modeled and Observed PCB Concentrations in Zooplankton, *Mysis*, and *Diporeia* at Saugatuck (1994 and 1995)



Agreement Between Modeled and Observed PCB Concentrations in Forage Fish at Saugatuck (1994 and 1995)



Agreement Between Modeled and Observed PCB Concentrations in Lake Trout at Saugatuck (1994 and 1995)

